

**ARIZONA GAME AND FISH DEPARTMENT  
HERITAGE DATA MANAGEMENT SYSTEM**

**Animal Abstract**

**Element Code:** AMACB02010  
**Data Sensitivity:** Yes

**CLASSIFICATION, NOMENCLATURE, DESCRIPTION, RANGE**

**NAME:** *Choeronycteris mexicana*  
**COMMON NAME:** Mexican Long-tongued Bat, Hog-nosed Bat  
**SYNONYMS:**  
**FAMILY:** Phyllostomidae

**AUTHOR, PLACE OF PUBLICATION:** Tschudi, J.J., 1844. Untersuchungen uber die Fauna Peruana. St. Gallen, parts 1-6, p. 72, 262pp.

**TYPE LOCALITY:** Mexico

**TYPE SPECIMEN:**

**TAXONOMIC UNIQUENESS:** In Arizona *Choeronycteris* is 1 of 3 genera in the family Phyllostomidae. *C. mexicana* is the only species in the genus that comes as far north as Arizona. *Choeronycteris mexicana* is regarded as a monotypic species by Koopman (in Wilson and Reeder 1993), and Simmons (in Wilson and Reeder 2005); nominal subspecies *ponsi* from northwestern Venezuela is now regarded as a subspecies of *Choeroniscus periosus* (Simmons, in Wilson and Reeder 2005). See Van Den Bussche (1992) for an analysis of phylogenetic relationships of phyllostomid bats based on restriction-site variation in the ribosomal-DNA gene complex.

**DESCRIPTION:** A rather large bat with a long, slender nose, and large eyes. It has a nose leaf that is broad at the base and pointed at the tip (looks like a small triangular bump near the nose tip), measuring about 5.0 mm high, which may help direct the ultrasonic echolocation signals the bat sends through its nostrils. Other measurements include the forearm between 42.0-48.0 mm (1.7-1.9 in.), a wingspan of 33-38 cm (13-15 in), the hind foot between 11.0-14.0 mm (0.44-0.56 in.), and the weights ranging between 10-25 grams. The tail is approximately 10 mm in length, about one-third the length of the naked interfemoral membrane. The dorsal pelage color varies from buffy brown to dark grayish-brown, palest on shoulders; venter is paler; ears pale brownish gray. The tongue is long and extendable, and can extend up to a third of their body length. Upper incisors are small, but do not fill the space between the canines. There are no permanent lower incisors, but one to four deciduous teeth may persist in adults. In flight, the wings make a swishing sound similar to that produced by long-nosed bats.

**AIDS TO IDENTIFICATION:** Species of the Phyllostomidae family found in Arizona, including *Choeronycteris mexicana*, are identified by the presence of a flap or leaf of skin extending from the tip of the nose. Bats of the other three families in Arizona lack such a

nasal leaf. *Leptonycteris curasoae yerbabuenae* lacks a visible tail and is larger (forearm 51.0-55.0 mm [2.04-2.2 in.]; hind foot 14.0-14.7 mm [0.56-0.59 in.]). The tail of *Macrotus* extends to slightly beyond the interfemoral membrane.

**ILLUSTRATIONS:**

B&W photo (Hoffmeister 1986:63)  
Color photo (Whitaker 1980: plate 180)  
Color photo (Wilson 1999)  
Color photo (Harvey 1999)  
Color photo (Whitaker 1996)

**TOTAL RANGE:** Southern California, southern Arizona, southwestern New Mexico, southern tip of Texas and much of northern and central Mexico. According to the Nevada Bat Working Group there was a single individual found in Las Vegas.

**RANGE WITHIN ARIZONA:** Southeast Arizona, from the Chiricahua Mountains extending as far north as the Santa Catalina Mountains and as far west as the Baboquivari Mountains. AGFD HDMS unpublished records show them in Pinal, Pima, Graham, Santa Cruz and Cochise counties.

**SPECIES BIOLOGY AND POPULATION TRENDS**

**BIOLOGY:** Peak activity for *Choeronycteris mexicana* occurs 1.5 hours after sunset and then at low levels until about 3 hours after sunset. They are less gregarious than other colonial bats and less inclined to roost with other bat species. In roosts, they do not cluster closely together but hang 2.0-5.0 cm (0.8-2.0 in.) apart, usually by only one foot so they can swivel 360° to detect predators. Roosts usually consist of 15 or fewer individuals, but when considering roosts in close proximity to each other, population numbers may reach up to 40-50. This species is thought to migrate seasonally to take advantage of suitable sources of food. They normally migrate south across the border into Mexico to spend the winter. Southern Arizona is at the extreme northern edge of its range, where it is found in sexually segregated and nursery colonies during the summer.

**REPRODUCTION:** Females segregate from the males, and according to Cockrum and Petryszyn (1992), only adult females move north from Mexico into the United States with the males remaining “in the southern part of the range during the time that young are being nourished by the mothers in the north.” The young (typically one baby) are born mid to late June and early July, but parturition may be as late as September in Mexico. As with many species of bats, the fetus is about 30% of the mother’s weight. Parturition usually lasts about 15 minutes, resulting in the birth of a neonate in a remarkably advanced state of development. The newborn bat is surprisingly well furred on the dorsum with a dense, dark pelage; the venter is scantily furred with silvery hair. Young grow rapidly and can probably fly within 2-

3 weeks. After the young become volant, these bats move about opportunistically in search of food. Females are known to carry their young in flight.

**FOOD HABITS:** This bat feeds on nectar, pollen, probably insects, and occasionally fruit of columnar cacti (these bats are not typically found in low desert situations). They especially feed on the flowers of paniculate agaves. The bristle-like tongue and lack of lower incisors, aid this bat in lapping up flower nectar and pollen.

During winter some are reported to feed at hummingbird feeders. It is not known if they are feeding on other things at that time.

**HABITAT:** Mesic areas in canyons of mixed oak-conifer forests in mountains rising from the desert; in Mexico includes arid thorn scrub, and tropical deciduous forests. Caves and abandoned mines are favored daytime retreats where they prefer to roost in the dimly lit areas often near the entrance. They are also often found in shallow caves or rock shelters. A few are found in palo verde-saguaro areas. Some of their range overlaps with *Leptonycteris*, but is not great (see Hevly 1979). *Choeronycteris* usually occupies higher elevations than *Leptonycteris* when it arrives in spring, and they may use the same roost year after year. Based on a study conducted by Carter and Peachey in 1996, all roost sites in the Cienega Creek Natural Preserve, except one, were located immediately adjacent to the creek. The roost sites consisted of pocketed, eroded clay soil holes such as sink holes, or soil piping caves. The majority of the soil piping caves were only a few meters long and 1-2 meters high, having a characteristic dome ceiling which seems to be where the bats prefer to roost.

**ELEVATION:** Records from 2,540 - 7,320 ft. (774 - 2,233 m), but most are from 4,000 - 6,000 ft. (1,220 - 1,830 m).

**PLANT COMMUNITY:** In the northern part of this bat's range, roost sites are commonly associated with mesic areas in oak-conifer woodlands or semi-desert grasslands. Dominant species include: oaks (*Quercus*), alligator juniper (*Juniperus deppeana*), manzanita (*Arctostaphylos*), yucca and agave. Near Tucson, they feed predominantly on cactus and *Agave* species. Cryan and Bogan (2003) observed species of *Agave* as the consistent floral characteristic of all sites visited, with *Agave schottii* observed blooming at occupied sites before mid-June, after which blooming *A. palmeri* was encountered.

**POPULATION TRENDS:** Unknown. Based on a recent study by Cryan and Bogan (2003), there is not sufficient evidence to conclude that populations in Arizona and New Mexico have increased or decreased in recent years. Searches of rock crevices and shelters in historical roost areas often revealed multiple roosting groups, suggesting that aggregations of *C. mexicana* are dispersed among several proximate sites.

**SPECIES PROTECTION AND CONSERVATION**

**ENDANGERED SPECIES ACT STATUS:** SC (USDI, FWS 1996)  
[C2 USDI, FWS 1989, 1991, 1994]

**STATE STATUS:** 2 (AZGFD, AWCS 2022)  
[1C (AGFD SWAP 2012)]  
[WSC (AGFD, WSCA 1996 in prep)]  
[Threatened (AGFD, TNW 1988)]

**OTHER STATUS:** Bureau of Land Management Sensitive  
(USDI, BLM AZ 2008, 2010, 2017)  
[Not Bureau of Land Management Sensitive  
(USDI, BLM AZ 2005)]  
[Bureau of Land Management Sensitive  
(USDI, BLM AZ 2000)]  
Forest Service Sensitive (USDA, FS Region  
3 2007, 2013)  
[Not Forest Service Sensitive (USDA, FS  
Region 3, 1999)]  
[Forest Service Sensitive USDA, FS Region  
3, 1988]  
NT (IUCN, Solari 2018)  
[LR/nt (Chiroptera Specialist Group 1996,  
*In:* IUCN 2006)]  
A, Determined Threatened (Secretaria de  
Medio Ambiente 2000, 2010)  
[Determined Threatened, Secretaria de  
Desarrollo Social 1994]

**MANAGEMENT FACTORS:** This species is very wary of humans and easily disturbed. They are difficult to survey for because they roost in small (5-15) colonies. Threats include recreational caving, mine reclamation, renewed mining, and loss of riparian habitat. Because of its propensity for moving between roosts in a small area, the loss of riparian vegetation may be a greater threat to the species than disturbance at a particular roost (Cryan and Bogan 2003). In addition, the loss of food resources (Agaves in Mexico [over harvesting]) due to development, fire or grazing may also have an affect on this species.

**PROTECTIVE MEASURES TAKEN:**

**SUGGESTED PROJECTS:** Restrict human disturbances at known roost sites. Due to their susceptibility to population decline, bat populations should be monitored. Research is needed on habitat needs, food habits, pollination role, survivorship, distribution, roosting patterns, and life history. In addition, studies looking at movement and revisits, possibly through banding, are needed, along with the possible affects of artificial feeders on their health.

**LAND MANAGEMENT/OWNERSHIP:** BIA - Tohono O'odham and San Carlos Reservations; BLM - Safford and Tucson Field Office; DOD - Fort Huachuca Military Reservation; NPS - Organ Pipe Cactus National Monument and Saguaro National Park; USFS - Coronado National Forest; State Land Department; Kartchner Caverns State Park; Pima County; Agua Caliente County Park; Cienega Creek Nature Preserve; AMNH Southwestern Research Station; TNC - Ramsey Canyon and Muleshoe Ranch Preserves; Private.

## SOURCES OF FURTHER INFORMATION

### REFERENCES:

- Arizona Game and Fish Department. 1988. Threatened native wildlife in Arizona. Arizona Game and Fish Department Publication. Phoenix, Arizona. p. 24.
- Arizona Game and Fish Department. 1996. Mammal Diversity Review notes.
- Arizona Game and Fish Department. 1996, in prep. Wildlife of special concern in Arizona. Arizona Game and Fish Department Publication. Phoenix, Arizona. p. 24.
- Arizona Game and Fish Department. 2012. Arizona's State Wildlife Action Plan 2012-2022. Arizona Game and Fish Department, Phoenix, Arizona. 233 pages.
- Arizona Game and Fish Department. 2022. Arizona Wildlife Conservation Strategy: 2022-2032. Arizona Game and Fish Department, Phoenix, Arizona. 378 pages.
- Arroyo-Cabrales, J., R.R. Hollander and J.K. Jones, Jr. 1987. *Choeronycteris mexicana* Tschudi, 1844, Mexican long-tongued bat. Mammalian Species 291:1-5. <https://doi.org/10.2307/3503823>
- Carter, D.E., Jr. and W.D. Peachey. 1996. Population study of Mexican long-tongued bats in southeast Pima County, Arizona. Heritage Grant I96024 report for Arizona Game and Fish Department, Phoenix Arizona. Pima County Parks and Recreation Department, Tucson, Arizona.
- Castner, S.V. 1995. Wildlife Views. Arizona Game and Fish Department. June 1995, p. 5.
- Chiroptera Specialist Group. 1996. *Choeronycteris mexicana*. In: IUCN 2006. 2006 IUCN Red List of Threatened Species. [www.iucnredlist.org](http://www.iucnredlist.org). Downloaded on 27 September 2006.
- Cockrum, E.L. and Y. Petryszyn. 1992. Seasonal and geographic distribution of the Mexican long-tongued bat *Choeronycteris mexicana* Tschudi. Draft.
- Constantine, D.G. 1987. Long-tongued bat and spotted bat at Las Vegas, Nevada. The Southwestern Naturalist 32(3):392. <https://doi.org/10.2307/3671459>
- Cryan, P.M. and M.A. Bogan. 2003. Recurrence of Mexican long-tongued bats (*Choeronycteris mexicana*) at historical sites in Arizona and New Mexico. Western North American Naturalist 63(3):314-319. <https://www.jstor.org/stable/41717299>
- Harvey, M.J., J.S. Altenbach, and T.L. Best. 1999. Bats of the United States. Arkansas Game and Fish Commission in cooperation with Ashville Field Office, U.S. Fish and Wildlife Service. p. 20.

- Hevly, R.H. 1979. Dietary habits of two nectar and pollen feeding bats in southern Arizona and northern Mexico. *Journal of the Arizona-Nevada Academy of Science* 14(1):13-18. <https://www.jstor.org/stable/40024466>
- Hoffmeister, D.F. 1986. *Mammals of Arizona*. University of Arizona Press, Tucson, Arizona and the Arizona Game and Fish Department, Phoenix, Arizona. pp.62-64.
- Huey, L.M. 1954. *Choeronycteris mexicana* from southern California and Baja California, Mexico. *Journal of Mammalogy* 35(3):436-437. <https://doi.org/10.2307/1375981>
- Jones, J.K., Jr., and D.C. Carter. 1976. Annotated checklist, with keys to subfamilies and genera. Pages 7-38 *In*: R.J. Baker, J.K. Jones, Jr., and D.C. Carter, editors. *Biology of bats of the New World family Phyllostomatidae. Part I. Special Publications The Museum Texas Tech University* 10:1-218.
- NatureServe Explorer: An online encyclopedia of life [web application]. 2001. Version 1.6. Arlington, Virginia, USA: NatureServe. Available: <https://explorer.natureserve.org/> (Accessed: 10 March 2003).
- Secretaría de Desarrollo Social. 1994. *Diario Oficial de la Federacion*. p. 25.
- Secretaría de Medio Ambiente. 2000. *Diario Oficial de la Federacion*. p. 43.
- Secretaría de Medio Ambiente y Recursos Naturales. 2010. NORMA Oficial Mexicana NOM-059-SEMARNAT-2010, Protección ambiental-Especies nativas de México de flora y fauna silvestres-Categorías de riesgo y especificaciones para su inclusión, exclusión o cambio-Lista de especies en riesgo.
- Solari, S. 2018. *Choeronycteris mexicana*. The IUCN Red List of Threatened Species 2018: e.T4776A22042479. <http://dx.doi.org/10.2305/IUCN.UK.2018-2.RLTS.T4776A22042479.en> (Accessed 21 February 2023)
- Turkowski, F.J. and A.W. Lewis. 1974. Distribution record of some Arizona mammals. *Journal of the Arizona Academy of Science* 9(3):89-90. <https://doi.org/10.2307/40023605>
- USDA, Forest Service, Region 3. 1988. Regional Forester's sensitive species list.
- USDA, Forest Service, Region 3. 1999. Regional Forester's sensitive species list.
- USDA, Forest Service, Region 3. 2007. Regional Forester's list of sensitive animals.
- USDA, Forest Service, Region 3. 2013. Regional Forester's list of sensitive animals.
- USDI, Bureau of Land Management. 2000. Arizona BLM sensitive species list. Instruction memorandum No. AZ-2000-018. Bureau of Land Management, Arizona State Office, Phoenix, Arizona.
- USDI, Bureau of Land Management. 2005. Arizona BLM sensitive species list. Bureau of Land Management, Arizona State Office, Phoenix, Arizona.
- USDI, Bureau of Land Management Region 2. 2008. Arizona BLM sensitive species list. Bureau of Land Management, Arizona State Office, Phoenix, Arizona.
- USDI, Bureau of Land Management Region 2. 2010. Arizona BLM sensitive species list. Instruction memorandum No. AZ-IM-2011-005. Bureau of Land Management, Arizona State Office, Phoenix, Arizona.
- USDI, Bureau of Land Management. 2017. Arizona BLM sensitive species list. Instruction memorandum No. AZ-IM-2017-009. Bureau of Land Management, Arizona State Office, Phoenix, Arizona. 6 pages.
- USDI, Fish and Wildlife Service. 1989. Endangered and threatened wildlife and plants; animal notice of review; notice of review. *Federal Register* 54(4):554-579.

- USDI, Fish and Wildlife Service. 1991. Endangered and threatened wildlife and plants; animal candidate review for listing as endangered or threatened species; notice of review. Federal Register 56(225):58804-5836.
- USDI, Fish and Wildlife Service. 1994. Endangered and Threatened Wildlife and Plants; Animal Candidate Review for Listing as Endangered or Threatened Species; Proposed Rule. Federal Register 59(219):58985.
- USDI, Fish and Wildlife Service. 1996. Endangered and threatened wildlife and plants; animal candidate review for listing as endangered or threatened species; notice of review. Federal Register 61(40):7596-7613.
- Van Den Bussche, R.A. 1992. Restriction-site variation and molecular systematics of New world leaf-nosed bats. Journal of Mammalogy 73(1):29-42.  
<https://doi.org/10.2307/1381863>
- Whitaker, J.O. 1980. The Audubon Society field guide to North American mammals. A.A. Knopf. New York, New York. p. 307; plate 180.
- Whitaker, J.O. 1996. The Audubon Society field guide to North American mammals. A.A. Knopf. New York New York. p. 318 plate 25.
- Wilson, D.E., and D.M. Reeder, editors. 1993. Mammal species of the world: a taxonomic and geographic reference. Second edition. Smithsonian Institution Press, Washington, DC. xviii + 1206 pages.
- Wilson, D.E., and D.M. Reeder, editors. 2005. Mammal species of the world: a taxonomic and geographic reference. Third edition. The Johns Hopkins University Press, Baltimore, Maryland. Two volumes. 2,142 pages. Available online at:  
<https://www.departments.bucknell.edu/biology/resources/msw3/>
- Wilson D.E. and S. Ruff, editors. 1999. The Smithsonian Book of North American Mammals. Smithsonian Institution Press, Washington, in association with the American Society of Mammalogists. pp. 75-76.

**MAJOR KNOWLEDGEABLE INDIVIDUALS:**

Dr. Paul Cryan – USGS, Fort Collins Science Center, Fort Collins, CO.  
Ronnie Sidner – Tucson, AZ.

**ADDITIONAL INFORMATION:**

In the 1980s (at least) in late August approximately 100 of these bats congregated in an old log homestead in Ramsey Canyon, Huachuca Mountains. Dr. E.L. Cockrum (pers comm 1992) speculated that the congregation may be due to the females and fledged young gathering near a food source, the numerous hummingbird feeders in the canyon.

Ronnie Sidner reports that they have been observing them more frequently at hummingbird feeders, and higher in the Santa Catalina Mountains.

*Choeronycteris mexicana*. From the Greek *choiros* meaning pig (refers to the pig-shaped snout) and *nykteris* meaning bat. The specific epithet, *mexicana*, refers to its major distribution, and where the species was first described.

**Revised:** 1991-12-26 (JSP)  
1992-05-03 (BKP)  
1992-09-28 (RBS)  
1994-03-29 (DCN)  
1997-03-03 (SMS)  
2003-03-10 (AMS)  
2006-09-27 (SMS)  
2023-02-21 (MBL)

To the user of this abstract: you may use the entire abstract or any part of it. We do request, however, that if you make use of this abstract in plans, reports, publications, etc. that you credit the Arizona Game and Fish Department. Please use the following citation:

Arizona Game and Fish Department. 20XX (= **year of last revision as indicated at end of abstract**). X...X (= **taxon of animal or plant**). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. X pp.