

**ARIZONA GAME AND FISH DEPARTMENT  
HERITAGE DATA MANAGEMENT SYSTEM**

**Animal Abstract**

**Element Code:** AMACD02020  
**Data Sensitivity:** Yes

**CLASSIFICATION, NOMENCLATURE, DESCRIPTION, RANGE**

**NAME:** *Eumops underwoodi*  
**COMMON NAME:** Underwood's Bonneted Bat, Underwood's Mastiff Bat  
**SYNONYMS:** *Eumops sonoriensis* Benson  
**FAMILY:** Molossidae

**AUTHOR, PLACE OF PUBLICATION:** Goodwin, 1940. Am. Mus. Novit., 1075:2.

**TYPE LOCALITY:** Honduras, La Paz, 6 km N. Chinacla, 3,000 feet.

**TYPE SPECIMEN:**

**TAXONOMIC UNIQUENESS:** In North America, *Eumops underwoodi* is 1 of 3 species in the genus *Eumops*; the others are *E. perotis* (SW United States), and *E. glaucinus* (southern tip of Florida). There are 2 subspecies recognized in *Eumops underwoodi*, *E. u. underwoodi* and *E. u. sonoriensis* (Hall 1981). *E. u. sonoriensis* is found in Arizona.

**DESCRIPTION:** Large bat, with total length 16.0-17.0 cm (6.4-6.8 in.), length of head and body about 13.0 cm; forearm 6.0-7.4 cm (2.6-2.96 in.); hind foot 1.5-1.8 cm (0.6-0.72 in.); weight 45-65 g. Long, narrow wings; wingspan 50.0-54.0 cm (20.0-21.6 in.). Ears 2.8-3.3 cm (1.12-1.32 in), meet at midline, projecting forward; ears just reach to tip of nose when laid forward; tragus small and rounded. Distal half of tail free from interfemoral membrane. Color varies from cinnamon to mummy brown and the under parts are hazel. Fur bicolored, lighter at base. A few long guard hairs on rump extend 7.0-10.0 mm (0.28-0.4 in.) beyond rest of fur. Males with well-developed gular gland (as all Molossidae). Skull short, wide, and unusually strongly ridged.

**AIDS TO IDENTIFICATION:** Superficially similar to all other free-tailed bats, but larger than *Tadarida*. Upper lip of *Tadarida* displays vertical wrinkles; lips of *Eumops* are smooth. *E. underwoodi* is most similar to *E. perotis* with which it may be sympatric in Arizona. *E. perotis* is slightly larger, forearm 7.3-8.3 cm (2.92-3.32 in.); and has much longer ears, 3.6-4.7 cm (1.44-1.88 in), reaching beyond nose when laid forward; tragus broad and square; and lacks long bristle-like hairs like those that protrude from rump of *E. underwoodi*. *E. u. underwoodi* color varies from a light yellowish to grayish-brown dorsally and is paler ventrally.

**ILLUSTRATIONS:**

Black and white photo (Barbour and Davis 1969: 228, 229, 230)  
Color photo (Barbour and Davis 1969: plate XXIII)  
Color photo (Whitaker 1980: plate 148)

Color photo (Wilson 1999)

Color photo (Harvey 1999)

**TOTAL RANGE:** Found from western Honduras through the western mainland of Mexico, reaching most northern part of its range just north of the US/Mexico border from Sasabe to Organ Pipe National Monument, Pima County, Arizona. However, according to Wilson (1999) and Hall (1981), the range is from southern Arizona to Sonora, Mexico.

**RANGE WITHIN ARIZONA:** Known from four localities near Sasabe and the vicinity of the Baboquivari Mountains, and from Organ Pipe Cactus National Monument, Pima County.

### **SPECIES BIOLOGY AND POPULATION TRENDS**

**BIOLOGY:** Despite their great size, and formidable dentition, these bats seem rather gentle. Their presence can be detected by their characteristic high pitched “peeps” emitted several times a minute when flying. These “peeps” are quite intense and may actually hurt the ears of an observer if the bat passes too close. Probably roosts in crevices along steep cliffs, perhaps in cracks of buildings. According to Cockrum (1981), the “day roosts are probably in high (30-40 feet or more above ground level) rock crevices in steep cliffs.” Narrow wings provide poor maneuverability; cannot take off from a flat surface. Water sites for drinking must provide a large open-water surface (15 to 30 feet or more) for the long gliding approach of this bat. It has been recorded at a flight speed of 43 km per hour.

Presumably a year-round resident in Arizona. Active during the warm months and hibernates in high rock crevices during winter. May occasionally be active during the winter. Cockrum (1981) reports male and female bats active at Organ Pipe Cactus National Monument during the months of January, March, May, June, August and September.

**REPRODUCTION:** Apparently, single young born in late June or July. Constantine (1961) reported a female gave birth on July 5 to a normal young which was successfully raised in captivity.

**FOOD HABITS:** Feeds on night flying insects 6 to 60 mm in length. Ross (1967) found *E. underwoodi* to take 13% short-horned grasshoppers and 47% scarab beetles. Very little is known about its natural feeding habits.

**HABITAT:** Their preferred habitat is not well understood. They have been netted over waterholes in desert and mesquite/grassland situations. In Arizona, they have been observed in Sonoran Desert habitat, while in Mexico in pine-oak forests (Pierson, Date unknown). While it is presumed that they roost primarily in rock crevices on cliff faces, the only identified roost was in a large, hollow tree in Jalisco, Mexico.

**ELEVATION:** 1,000 ft. (305 m) at Organ Pipe Cactus National Monument, 4,000 ft. (1,220 m) near Sasabe. According to AGFD HDMS unpublished records (accessed 2003), the elevation ranges from 1,080 – 3,760 ft (329 – 1,146 m).

**PLANT COMMUNITY:** Sonoran Desert Scrub; mesquite/grassland. Associated vegetation includes: mesquite, willow, saguaro, ocotillo, cholla, and prickly pear.

**POPULATION TRENDS:** Unknown.

### **SPECIES PROTECTION AND CONSERVATION**

**ENDANGERED SPECIES ACT STATUS:** SC (USDI, FWS 1996)  
[C2 (USDI, FWS 1985, 1989, 1991, 1994)]  
[C2 (USDI, FWS 1982) at subspecies *E. u. sonoriensis*]

**STATE STATUS:** 2 (AZGFD, AWCS 2022)  
[1B (AGFD SWAP 2012)]

**OTHER STATUS:** Not Forest Service Sensitive (USDA, FS Region 3 1999, 2007, 2013)  
[Forest Service Sensitive USDA, FS Region 3, 1988]  
Not Bureau of Land Management Sensitive (USDI, BLM 2008, 2010, 2017)  
[Bureau of Land Management Sensitive (USDI, BLM AZ 2000, 2005)]  
LC (IUCN, Miller et al. 2016)

**MANAGEMENT FACTORS:** Assuming that this species is primarily cliff dwelling, it could be threatened by any activities that destroy cliff habitat (recreational climbing, water impoundments, highway construction, quarry operations). Grazing and pesticide applications in agricultural areas could impact foraging habitat.

### **PROTECTIVE MEASURES TAKEN:**

**SUGGESTED PROJECTS:** More surveys are needed to delineate the range of this species in the southwestern U.S. In addition, surveys and studies are needed on distribution of breeding colonies, seasonal movements, roosting and foraging requirements, and life history. Methods need to be described to determine whether this species can be distinguished from other large molossid acoustically.

**LAND MANAGEMENT/OWNERSHIP:** BIA - Tohono O'odham Nation; FWS - Buenos Aires National Wildlife Refuge; NPS - Organ Pipe Cactus National Monument; Private.

**SOURCES OF FURTHER INFORMATION****REFERENCES:**

- Arizona Game and Fish Department. 1996. Mammal Diversity Review notes. Arizona Game and Fish Department, Phoenix, Arizona.
- Arizona Game and Fish Department. 2012. Arizona's State Wildlife Action Plan 2012-2022. Arizona Game and Fish Department, Phoenix, Arizona. 233 pages
- Arizona Game and Fish Department. 2022. Arizona Wildlife Conservation Strategy: 2022-2032. Arizona Game and Fish Department, Phoenix, Arizona. 378 pages.
- Barbour, R.W. and W.H. Davis. 1969. Bats of America. The University Press of Kentucky. pp. 228-231.
- Cockrum, E.L. 1960. The recent mammals of Arizona: their taxonomy and distribution. The University of Arizona Press. Tucson. pp. 66-67.
- Cockrum, E.L. 1981. Bat populations and habitats at the Organ Pipe Cactus National Monument. Technical report No. 7. Cooperative National Park Resources Studies Unit, University of Arizona, Tucson, Arizona. p. 31.
- Cockrum, E.L. and A.L. Gardner. 1960. Underwood's mastiff bat in Arizona. Journal of Mammalogy 41(4):510-511. <https://doi.org/10.2307/1377544>
- Constantine, D.G. 1961. Locality records and notes on western bats. Journal of Mammalogy 42(3):404-405. <https://doi.org/10.2307/1377050>
- Eger, J.L. 1977. Systematics of the genus *Eumops* (Chiroptera: Molossidae). Royal Ontario Museum. Life Sciences Contributions 110:1-69.
- Goodwin, G.G. 1940. Three new bats from Honduras and the first record of *Enchisthenes harti* (Thomas) for North America. American Museum Novitates 1075:1-3.
- Hall, E.R. 1981. The mammals of North America. Second Edition. John Wiley & Sons. New York, New York. p. 249.
- Harvey, M.J, J.S. Altenbach, and T.L. Best. 1999. Bats of the United States. Arkansas Game and Fish Commission in cooperation with Ashville Field Office, U.S. Fish and Wildlife Service. p. 58.
- Hoffmeister, D.F. 1986. Mammals of Arizona. The University of Arizona Press, Tucson, Arizona and The Arizona Game and Fish Department, Phoenix, Arizona. pp. 122-124.
- Miller, B., F. Reid, J. Arroyo-Cabrales, A.D. Cuarón, and P.C. de Grammont. 2016. *Eumops underwoodi*. The IUCN Red List of Threatened Species 2016: e.T8248A22025754. <http://dx.doi.org/10.2305/IUCN.UK.2016-2.RLTS.T8248A22025754.en>
- NatureServe Explorer: An online encyclopedia of life [web application]. 2001. Version 1.6. Arlington, Virginia, USA: NatureServe. Available: <https://explorer.natureserve.org/> (Accessed: 2 April 2003).
- Pierson, E.D. Date unknown. *Eumops underwoodi*. Species account.
- Petryszyn, Y. and E.L. Cockrum. 1990. Mammals of the Quitobaquito management area, Organ Pipe Cactus National Monument, Arizona. Technical report No. 36. Cooperative National Park Resources Studies Unit, University of Arizona, Tucson, Arizona. 32 pp.
- Ross, A. 1967. Ecological aspects of the food habits of insectivorous bats. Proceedings of the Western Foundation of Vertebrate Zoology 1:203-264.
- USDA, Forest Service Region 3. 1988. Regional Forester's sensitive species, Region 3, August 1988. U.S. Forest Service. 41 pages.

- USDA, Forest Service Region 3. 1999. Regional Forester's list of sensitive animals - 7/21/1999. U.S. Forest Service. 7 pages.
- USDA, Forest Service Region 3. 2000. Regional Forester's sensitive species list (7/21/99 as corrected 2/23/00). U.S. Forest Service. 13 pages.
- USDA, Forest Service Region 3. 2007. Regional Forester's list of sensitive animals. U.S. Forest Service. 39 pages.
- USDA, Forest Service Region 3. 2013. Regional Forester's sensitive species: animals - 2013. U.S. Forest Service. 5 pages.
- USDI, Bureau of Land Management. 2000. Arizona BLM sensitive species list. Instruction memorandum No. AZ-2000-018. Bureau of Land Management, Arizona State Office, Phoenix, Arizona.
- USDI, Bureau of Land Management. 2005. Arizona BLM sensitive species list. Bureau of Land Management, Arizona State Office, Phoenix, Arizona.
- USDI, Bureau of Land Management Region 2. 2008. Arizona BLM sensitive species list. Bureau of Land Management, Arizona State Office, Phoenix, Arizona.
- USDI, Bureau of Land Management Region 2. 2010. Arizona BLM sensitive species list. Instruction memorandum No. AZ-IM-2011-005. Bureau of Land Management, Arizona State Office, Phoenix, Arizona.
- USDI, Bureau of Land Management. 2017. Arizona BLM sensitive species list. Instruction memorandum No. AZ-IM-2017-009. Bureau of Land Management, Arizona State Office, Phoenix, Arizona. 6 pages.
- USDI, Fish and Wildlife Service. 1982. Endangered and threatened wildlife and plants; review of vertebrate wildlife for listing as endangered or threatened species; notice of review. Federal Register 47(251):58454-58460.
- USDI, Fish and Wildlife Service. 1985. Endangered and threatened wildlife and plants; review of vertebrate wildlife; notice of review. Federal Register 50(181):37958-37967.
- USDI, Fish and Wildlife Service. 1989. Endangered and threatened wildlife and plants; animal notice of review; notice of review. Federal Register 54(4):554-579.
- USDI, Fish and Wildlife Service. 1991. Endangered and threatened wildlife and plants; animal candidate review for listing as endangered or threatened species; notice of review. Federal Register 56(225):58804-58836.
- USDI, Fish and Wildlife Service. 1994. Endangered and threatened wildlife and plants; animal candidate review for listing as endangered or threatened species; notice of review. Federal Register 59(219):58982-59028.
- USDI, Fish and Wildlife Service. 1996. Endangered and threatened wildlife and plants; review of plant and animal taxa that are candidates for listing as endangered or threatened species; notice of review. Federal Register 61(40):7596-7613.
- Whitaker, J.O. 1980. The Audubon Society field guide to North American mammals. A.A. Knopf. New York, New York. pp. 335-336; plate 148.
- Wilson D.E. and S. Ruff, editors. 1999. The Smithsonian Book of North American Mammals. Smithsonian Institution Press, Washington, D.C., in association with the American Society of Mammalogists. pp. 134-135.

**MAJOR KNOWLEDGEABLE INDIVIDUALS:**

Yar Petryszyn - University of Arizona, Tucson.

**ADDITIONAL INFORMATION:**

No diurnal roost sites not known in Arizona.

Constantine (1961) found these bats were amenable to captivity after he trained them to eat mealworms. The one born in captivity was remarkably tame, squeaking to attract attention, scampering across the cage to meet visitors, jumping to a hand and running up an arm whenever it had the opportunity, and squeaking when company left.

Ronnie Sidner reported that the original site that they were taken, has disappeared, but they have been caught nearby. They are mainly found along large bodies of water (AGFD 1996).

**Revised:** 1991-01-08 (JSP)  
1992-05-26 (BKP)  
1992-10-02 (RBS)  
1994-03-25 (DCN)  
1996-06-19 (SMS)  
1997-03-04 (SMS)  
2003-04-13 (AMS)  
2023-02-27 (MBL)

To the user of this abstract: you may use the entire abstract or any part of it. We do request, however, that if you make use of this abstract in plans, reports, publications, etc. that you credit the Arizona Game and Fish Department. Please use the following citation:

Arizona Game and Fish Department. 20XX (= **year of last revision as indicated at end of abstract**). X...X (= **taxon of animal or plant**). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. X pp.